



Ingredients

Each vegetable capsule contains:

Cranberry concentrate

(*Vaccinium macrocarpon aiton (Ericaceae)*)



600 mg of 107x concentrate equal to 64,200 mg

Extracted from 100% whole ripe red cranberries.

Cranberry

A healthy
urinary tract starts with
CRANBERRY 107x

PREVENT AND
FIGHT INFECTIONS
THE NATURAL WAY.



CRANBERRY 107x
Sold exclusively to finer health food stores
www.newrootsherbal.com/store



CRANBERRY 107x treats urinary tract infections by taking advantage of the cranberry's legendary antibacterial properties. The special proanthocyanidin extract we have created delivers the benefits of 64,200 mg of fresh, whole, ripe cranberries in every capsule. As a result, you can trust **CRANBERRY 107x** to be fast-acting, with consistent results every time.

A natural remedy.

Modern medical research has revealed the chemical and physiological effects cranberries have on the urinary tract and how supplementing cranberry concentrates can help prevent urinary tract infections.

OUR CRANBERRY
CONCENTRATE OFFERS
THE MOST POTENT,
SPECIALLY STRUCTURED
PROANTHOCYANIDINS
AVAILABLE.

Why CRANBERRY Works.

Cranberries contain specially structured proanthocyanidins (PACs) that inhibit the adhesion of bacteria to the urinary tract and bladder, thereby preventing them from reproducing. Because of this, PACs are pivotal in the prevention of urinary tract infections.

There's more to CRANBERRY.

While cranberries are best known for their effect on urinary tract health, recent research indicates that they may also act against bacteria elsewhere in the body. The presence of cranberry has been shown to inhibit the adhesion and subsequent infection from many types of bacteria, including those that cause stomach ulcers and periodontal gum disease.

Suggested Use

To Prevent:

Take orally 2 capsules once or twice a day with water before meals, or as directed by your health care practitioner.

For a Urinary Tract Infection:

You must "catch" the infection immediately. Take orally 4 capsules with water before meals, 3 times daily, for 5 days. Drink 8 glasses of water daily.

If there is no relief from symptoms after 2 days, see your health care practitioner immediately.

References:

1. Ahuja, S., Kaack, B. and Roberts, J. (1998).
2. Allison, D.G., *et al.* (2000).
3. Avorn, J. (1996).
4. Avorn, J., *et al.* (1994).
5. Burger, O., Weiss, E., Sharon, N., Tabak, M., Neeman, I., and Ofek, I. (2002).
6. Foo, Y.P., Lu, Y., Howell, A.B. and Vorsa, N. (2000).
7. Habash, M.B., *et al.* (1999).
8. Harkins, K.J. (2000).
9. Howell, A.B., Leahy, M., Kurowska, E., and Guthrie, N. (2001).
10. Jepson, R.G., Hihaljevic, L., and Craig, J. (2000).

